

The SPICT™ is a guide to identifying people at risk of dying within the next 12 months.

## Look for two or more general indicators of deteriorating health.

- Performance status poor or deteriorating, with limited reversibility. (needs help with personal care, in bed or chair for 50% or more of the day).
- Two or more unplanned hospital admissions in the past 6 months.
- Weight loss (5 - 10%) over the past 3 - 6 months and/or body mass index < 20.
- Persistent, troublesome symptoms despite optimal treatment of any underlying condition(s).
- Lives in a nursing care home or NHS continuing care unit, or needs care to remain at home.
- Patient requests supportive and palliative care, or treatment withdrawal.

## Look for any clinical indicators of advanced conditions

### Cancer

Functional ability deteriorating due to progressive metastatic cancer.

Too frail for oncology treatment or treatment is for symptom control.

### Dementia/ frailty

Unable to dress, walk or eat without help.

Choosing to eat and drink less; difficulty maintaining nutrition.

Urinary and faecal incontinence.

Unable to communicate meaningfully; little social interaction.

Fractured femur; multiple falls.

Recurrent febrile episodes or infections; aspiration pneumonia.

### Neurological disease

Progressive deterioration in physical and/or cognitive function despite optimal therapy.

Speech problems with increasing difficulty communicating and/or progressive dysphagia.

Recurrent aspiration pneumonia; breathless or respiratory failure.

### Heart/ vascular disease

NYHA Class III/IV heart failure, or extensive, untreatable coronary artery disease with:

- breathlessness or chest pain at rest or on minimal exertion.

Severe, inoperable peripheral vascular disease.

### Respiratory disease

Severe chronic lung disease with:

- breathlessness at rest or on minimal exertion between exacerbations.

Needs long term oxygen therapy.

Has needed ventilation for respiratory failure or ventilation is contraindicated.

### Kidney disease

Stage 4 or 5 chronic kidney disease (eGFR < 30ml/min) with deteriorating health.

Kidney failure complicating other life limiting conditions or treatments.

Stopping dialysis.

### Liver disease

Advanced cirrhosis with one or more complications in past year:

- diuretic resistant ascites
- hepatic encephalopathy
- hepatorenal syndrome
- bacterial peritonitis
- recurrent variceal bleeds

Liver transplant is contraindicated.

## Assess and plan supportive & palliative care

- Review current treatment and medication so the patient receives optimal care.
- Consider referral for specialist assessment if symptoms or needs are complex and difficult to manage.
- Agree current and future care goals/ plan with the patient and family.
- Plan ahead if the patient is at risk of loss of capacity.
- Handover: care plan, agreed levels of intervention, CPR status.
- Coordinate care (eg. with a primary care register).